

## Gemini Series Isotoma

*I. hybrida*

Approximate seed count (tuned): 280,000-340,000 sds/oz, 10000 – 12000 sds/gr.

### Plug Production

#### Media

Use a well-drained, disease-free, media with a pH of 5.5 to 6.2 and an EC of 0.75 mmhos/cm.

#### Sowing

Sow 2 to 4 seeds per cell in 288 or larger plug tray; no cover needed. Spray preventively with fungicide.

**Stage 1** – Germination begins between day 8 to 12 continuing through day 14 to 21.

**Soil temperature:** 68 to 72°F (20 to 22°C)

**Light:** Not necessary but beneficial

**Moisture:** Keep soil moist (level 4) in Stage 1.

**Humidity:** Maintain 95 to 97% relative humidity (RH) until cotyledons emerge.

#### Stage 2

**Soil temperature:** 68 to 72°F (20 to 22°C)

**Light:** Can be up to 2,500 f.c. (26,900 Lux).

**Moisture:** Maintain soil moisture moist (level 4), to allow the roots to penetrate into the media, don't let the media dry out.

**Fertilizer:** Apply fertilizer at rate 1 (less than 100 ppm Nitrate/0.2 – 0.5 mmhos/cm EC) from nitrate-form fertilizers.

#### Stage 3

**Soil temperature:** 65 to 68°F (18 to 20°C)

**Light:** Can be up to 2,500 f.c. (26,900 Lux). In low light area, supplemental light of 150 f.c. (1620 lux) is beneficial for reducing seedling stretching and cutting plug crop time.

**Moisture:** Maintain the moisture level constantly medium moist to medium wet (level 3-4)

**Fertilizer:** Apply fertilizer at rate 1 (less than 100 ppm Nitrate/less than 0.7 mmhos/cm EC) from nitrate-form fertilizers.

**Growth regulators:** None

#### Stage 4

**Soil temperature:** 65 to 68°F (18 to 20°C)

**Light:** can be up to 5,000 f.c. (53,800 Lux) if temperatures can be maintained.

**Moisture:** Same as Stage 3.

**Fertilizer:** Same as Stage 3.

### Growing On to Finish

#### Media

Use a well-drained, disease-free, soilless media with a pH of 5.5 to 6.2 and an EC of 0.75 mmhos/cm).

#### Temperature

**Nights:** 54 to 57°F (12 to 14°C)

**Days:** 60 to 65°F (16 to 18°C)

#### Light

Maintain light levels as high as possible while maintain moderate temperature

#### Photoperiod

Isotoma flowers most rapidly and uniformly at a minimum of 13 to 14 hours daylength.

#### Irrigation

Maintain media constantly moisture (level 2-3). Avoid plants dry to wilt.

#### Fertilizer

Gemini Isotoma is a moderate feeder. Apply fertilizer at rate 3 (175 to 225 ppm N/1.2 to 1.5 mS/cm) weekly or as needed with predominantly nitrate-form fertilizer with low phosphorus and high potassium, such as 15-5-15.

#### Growth Regulators

Isotoma is responsive to B-Nine/Alar (daminozide) 2500 – 3250 ppm (3.0-3.9 g/l of 85% formulation or 4.0 –5.0 g/l 64% formulation) at 2 weeks after transplant. If necessary, repeat the treatment a week later.

#### Pinching

Do not pinch.

### Crop Scheduling

**Sow to transplant (288 cell plug):** 4 to 5 weeks

**Transplant to flower:**

Container Size	Plants Per Pot	Weeks From Transplant	Total Weeks
4-in. (10.5-cm)	1	9 to 11	13 to 16
5-in. (13-cm)	1	10 to 12	14 to 17
Gallon (19-cm)	3	10 to 11	14 to 16

Under proper day length and temperature ranges

**Common Problems**

**Insect:** Aphid, Thrips, Spider mite, Sciara (in young plant stage)

**Disease:** Pythium, Bothrytis

**Note:**

Growers should use the information presented here as a starting point. Crop times will vary depending on the climate, location, time of year, and greenhouse environmental conditions. Chemical and PGR recommendations are only guidelines. It is the responsibility of the applicator to read and follow all the current label directions for the specific chemical being used in accordance with all regulations

630 231-1400  
panamseed.com

© 2011 Ball Horticultural Company 12109 REV 12/11  
™ denotes a trademark of and ® denotes a registered trademark of Ball Horticultural Company in the U.S. It may also be registered in other countries.

PanAmerican Seed.